


1 / 1 0

FIG.1
GPC PEAK PROFILE OF PRODUCT IN EXAMPLE 1



2 / 10





FIG. 2 ^1H -NMR SPECTRUM OF PRODUCT. IN EXAMPLE 1

3 / 10

FIG. 3

GPC. PEAK PROFILE OF PRODUCT IN EXAMPLE 2



4 / 10





FIG. 4 ^1H -NMR SPECTRUM OF PRODUCT IN EXAMPLE 2

5 / 10

FIG.5


GPC PEAK PROFILE OF PRODUCT IN EXAMPLE 3



6 / 10

FIG. 6


¹H-NMR SPECTRUM OF PRODUCT IN EXAMPLE 3



7 / 10

FIG. 7


^1H -NMR SPECTRUM OF PRODUCT IN EXAMPLE 4



8 / 10

FIG. 8

^1H -NMR SPECTRUM OF PRODUCT IN COMPARATIVE EXAMPLE 1




9 / 1 0

FIG. 9

PLOT OF CHANGE WITH TIME IN RESIDUAL ACRYLIC ACID AS COMPARISON BETWEEN
SYNTHESES IN EXAMPLE 4 AND COMPARATIVE EXAMPLE 1

ACRYLIC ACID CONVERSION*¹



*1 GPC ANALYZED DATA

*2 CALCULATED VALUE FROM INTEGRAL OF ¹H-NMR SPECTRUM OF
PRODUCT AFTER REMOVAL OF LOW BOILING COMPONENTS

10 / 10

FIG.10

COMPARISON IN GPC PEAK PROFILE BETWEEN PRODUCT IN EXAMPLE 4
AND COMPARATIVE EXAMPLE 1

